

GE Discovery LS 16 Body CT

ORAL CONTRAST PREPARATION INSTRUCTIONS

Type of Oral Contrast	None	Negative (H2O)	Positive (Readi-Cat or *Gastroview)	Special (VoLumen)
CT Protocol	Pelvic CT (fracture) CT Urogram (NC) CTA Abd/Pel/LE CTA Endograft CT Venogram Chest CT CTA Chest HR Chest CT	Liver CT Adrenal CT CT Urogram (with)	Abdomen CT Pelvic CT Abd/Pel CT Chest/Abd/Pel CT	Pancreas CT CT Enterography
Instructions (Abd ONLY)		<ul style="list-style-type: none"> 16 oz. H2O immediately before test (CT technologist will advise) 	<p>1.3 Bottles</p> <ul style="list-style-type: none"> 1 Bottle 1 hr before test 1/3 bottle immediately before test (CT technologist will advise) 	<p>Pancreas:</p> <p>1 Bottle</p> <ul style="list-style-type: none"> 1 Bottle VoLumen 1 hr prior to exam. 8 oz. H2O 15 min prior to exam
Instructions (Abd/Pel)		<ul style="list-style-type: none"> 16 oz. H2O immediately before test (CT technologist will advise) 	<p>2 Bottles</p> <ul style="list-style-type: none"> 1 Bottle 3 hrs before test 2/3 Bottle 30 min before test 1/3 bottle immediately before test (CT technologist will advise) 	<p>Enterography:</p> <p>2 Bottles</p> <ul style="list-style-type: none"> 2 Bottles VoLumen 1 hr before exam 16 oz. H2O immediately before test (CT technologist will advise)

*Gastroview: In case of suspected bowel perforation (trauma, diverticulitis, appendicitis, etc) use Gastroview INSTEAD of Readi-Cat. If the patient is at high risk for aspiration, DO NOT use Gastroview. Consult the Radiologist as needed.